



Laser speckle contrast imaging: age-related changes in microvascular blood flow and correlation with pulse-wave velocity in healthy subjects

Submitted by Anne Humeau-Heurtier on Wed, 11/26/2014 - 07:03

Titre	Laser speckle contrast imaging: age-related changes in microvascular blood flow and correlation with pulse-wave velocity in healthy subjects
Type de publication	Article de revue
Auteur	Khalil, Adil [1], Humeau-Heurtier, Anne [2], Mahé, Guillaume [3], Abraham, Pierre [4]
Editeur	Society of Photo-optical Instrumentation Engineers
Type	Article scientifique dans une revue à comité de lecture
Année	2015
Langue	Anglais
Date	Jan-05-2015
Numéro	5
Pagination	article id. 051010
Volume	20
Titre de la revue	Journal of Biomedical Optics
ISSN	1083-3668
Mots-clés	Blood [5], Blood flow [6], Laser speckle contrast imaging [7]
Résumé en anglais	<p>In the cardiovascular system, the macrocirculation and microcirculation—two subsystems—can be affected by aging. Laser speckle contrast imaging (LSCI) is an emerging noninvasive optical technique that allows the monitoring of microvascular function and can help, using specific data processing, to understand the relationship between the subsystems. Using LSCI, the goals of this study are: (i) to assess the aging effect over microvascular parameters (perfusion and moving blood cells velocity, MBCV) and macrocirculation parameters (pulse-wave velocity, PWV) and (ii) to study the relationship between these parameters. In 16 healthy subjects (20 to 62 years old), perfusion and MBCV computed from LSCI are studied in three physiological states: rest, vascular occlusion, and post-occlusive reactive hyperaemia (PORH). MBCV is computed from a model of velocity distribution. During PORH, the experimental results show a relationship between perfusion and age ($R^2=0.67$) and between MBCV and age ($R^2=0.72$), as well as between PWV and age at rest ($R^2=0.91$). A relationship is also found between perfusion and MBCV for all physiological states ($R^2=0.98$). Relationships between microcirculation and macrocirculation (perfusion-PWV or MBCV-PWV) are found only during PORH with $R^2=0.76$ and $R^2=0.77$, respectively. This approach may prove useful for investigating dysregulation in blood flow.</p>
URL de la notice	http://okina.univ-angers.fr/publications/ua5533 [8]
DOI	10.1117/1.JBO.20.5.051010 [9]

Lien vers le document <http://dx.doi.org/10.1117/1.JBO.20.5.051010> [9]
Titre abrégé J. Biomed. Opt

Liens

- [1] <http://okina.univ-angers.fr/akhalil/publications>
- [2] <http://okina.univ-angers.fr/a.hum/publications>
- [3] [http://okina.univ-angers.fr/publications?f\[author\]=1153](http://okina.univ-angers.fr/publications?f[author]=1153)
- [4] <http://okina.univ-angers.fr/pierre.abraham/publications>
- [5] [http://okina.univ-angers.fr/publications?f\[keyword\]=1795](http://okina.univ-angers.fr/publications?f[keyword]=1795)
- [6] [http://okina.univ-angers.fr/publications?f\[keyword\]=10142](http://okina.univ-angers.fr/publications?f[keyword]=10142)
- [7] [http://okina.univ-angers.fr/publications?f\[keyword\]=10143](http://okina.univ-angers.fr/publications?f[keyword]=10143)
- [8] <http://okina.univ-angers.fr/publications/ua5533>
- [9] <http://dx.doi.org/10.1117/1.JBO.20.5.051010>

Publié sur *Okina* (<http://okina.univ-angers.fr>)